

## Neurocirugía



https://www.revistaneurocirugia.com

## C0346 - INTRACRANIAL DYNAMICS IN THE SUPINE AND STANDING POSITIONS: DOES ANYTHING MORE CHANGE THAN THE MEAN INTRACRANIAL PRESSURE?

**P. López Gómez**, J. Esteban García, G. García Catalán, C. Bucheli Peñafiel, I. Valduvieco Juaristi, I. Pinto Rafael and R. Martín Láez

Hospital Universitario Marqués de Valdecilla, Cantabria, Spain.

## Resumen

**Objectives:** Little is known about variation in dynamics of intracranial pressure (ICP) when noncritically ill patients change from supine to standing position. Literature is scarce, most of it limited to modification of absolute ICP values. The aim of this study is to determine the variation of parameters extracted from ICP monitoring during changes from supine to standing position.

**Methods:** We retrospectively reviewed patients that underwent computerized ICP monitoring in whom at least 30 minutes of data before and after a postural change from supine to standing was available. Non-shunted patients with normal quantitative and qualitative nocturnal ICP recording were selected for analyses. Raw ICP data was automatically analyzed using ICP Digital Tools<sup>®</sup>, extracting the following parameters for both body positions: mean ICP, dispersion, cerebral pulse pressure (PP), systolic raising coefficient (dPdt), pressure-pulse correlation coefficient (RAP), partial pressure-pulse regression coefficient (RPPC) and closing hydrostatic pressure (P<sub>0</sub>). SPSS 21 was used for statistical analysis, considering significance at p < 0.05.

**Results:** Of 308 patients studied, 217 have monitoring during positional changes. In 31 of those cases, overnight ICP recording was considered to be quantitative and qualitatively normal. In those subjects mean ICP decreased from  $3.66 \pm 3.17$  mmHg in supine to  $-5.62 \pm 3.63$  mmHg in standing (p < 0.0001) and P<sub>0</sub> from  $-1.74 \pm 13.60$  to  $-15.83 \pm 10.15$  mmHg (p < 0.0001). Dispersion increased from  $1.79 \pm 0.94$  to  $2.20 \pm 0.98$  mmHg (p = 0.008), PP from  $2.88 \pm 0.93$  to  $3.58 \pm 1.24$  mmHg (p < 0.0001) and dPdt from  $15.15 \pm 6.64$  to  $17.79 \pm 7.17$  mmHg/s (p = 0.006). No statistically significant changes were detected in RAP nor in RPPC.

**Conclusions:** ICP variations during positional changes are not limited to mean ICP decrease, as P0 decreases as well. Oppositely, ICP dispersion, PP and dPdt rise significantly.