



Neurocirugía



<https://www.revistaneurocirugia.com>

C0495 - USE OF ANTI-D WAVE TO OPTIMIZE AND MAXIMIZE SAFE RESECTION OF SPINAL CORD TUMORS, WHILE PRESERVING NEUROLOGICAL FUNCTIONS

L.H. González-Quarante, A. Traba López, R. González Rodrigálvarez, Ó. Lucas Gil de Sagredo del Corral, A. Montalvo Afonso, C. Fernández Carballal and R. García Leal

Hospital General Universitario Gregorio Marañón, Madrid, Spain.

Resumen

Objectives: To present the novel “anti D wave”, as a new neurophysiological tool to perform intraoperative neurophysiological monitoring in spinal cord tumors. We present our experience with an illustrative case of intraspinal ependymoma affecting the cervical cord.

Methods: We performed a review of the literature and found one thorough paper presenting the use of “anti D wave” or antidromic wave in human patients with tumoral diseases. This wave is deemed to be presumably used in a similar fashion than the better known “D wave”. Therefore, we systematically started to complement our standard intraoperative neurophysiological monitoring with the Antidromic Cortico Spinal Tract Potential (“anti-D” wave) in patients with spinal cord tumors.

Results: We present the video of an illustrative case of a 44-year-old male patient who presented at our institution complaining of a 6-month history of cervicalgia. MRI showed an intradural intramedullary cystic lesion that extended from C2 to C6. A small enhancing nodule was observed in the middle of the lesion. Surgical procedure with intraoperative neurophysiological monitoring was scheduled. In addition to the standard monitoring [somatosensory evoked potentials, transcranially elicited muscle motor evoked potentials, epidural motor evoked potentials (D wave) and cortical spinal cord-evoked potentials], we recorded the Antidromic Cortico Spinal tract Potential, also known as Anti-D wave.

Conclusions: We found a close correlation between anti-D wave and D-wave. We have recorded the anti-D wave in 3 additional patients with spinal cord tumors, observing a similar positive predictive value for preservation of neurological functions, when comparing this novel wave to the more commonly used D wave. Admittedly, further cases are needed to better characterize this new neurophysiological tool and add it to the intraoperative armamentarium for treating complex spinal cord tumors, yielding a better chance of preserving neurological functions.