

## Neurocirugía



https://www.revistaneurocirugia.com

## O-039 - ANEURYSMAL SUBARACHNOID HAEMORRHAGE: VOLUMETRIC QUANTIFICATION OF THE BLOOD DISTRIBUTION PATTERN TO ACCURATELY PREDICT THE RUPTURED ANEURYSM LOCATION

A. Mosteiro<sup>1</sup>, D. Culebras<sup>1</sup>, D. Vargas<sup>1</sup>, J.L. Moreno<sup>1</sup>, A. López-Rueda<sup>1</sup>, L. Llull<sup>1</sup>, D. Santana<sup>1</sup>, L. Pedrosa<sup>2</sup>, S. Amaro<sup>1</sup>, R. Torné<sup>1</sup> and J. Enseñat<sup>1</sup>

<sup>1</sup>Hospital Clínic de Barcelona, Barcelona, Spain; <sup>2</sup>Instituto de Investigaciones Biomédicas August Pi i Sunyer, Barcelona, Spain.

## Resumen

**Introduction:** In spontaneous subarachnoid haemorrhage (SAH) accurate determination of the bleeding source is paramount to guide treatment. Traditionally, the bleeding pattern has been used to predict the aneurysm location.

**Objectives:** Here, we have tested a software-based tool, which quantifies the volume of intracranial blood and stratifies it according to the regional distribution, to predict the location of the ruptured aneurysm.

**Methods:** A consecutive series of SAH patients admitted to a single tertiary centre between 2012 and 2018, within 72h of onset, harbouring a single intracranial aneurysm. A semi-automatized method of blood quantification, based on the relative density increase, was applied to initial non-contrast CTs. Five regions were used to define the bleeding patterns and to correlate them with aneurysm location: perimesencephalic, interhemispheric, right/left hemisphere and intraventricular.

**Results:** 68 patients were included for analysis. There was a strong association between the distribution of blood and the aneurysm location (p < 0.001). In particular: ACom and interhemispheric fissure (p < 0.001), MCA and ipsilateral hemisphere (p < 0.001), ICA and ipsilateral hemisphere and perimesencephalic cisterns (p < 0.001), PCom and hemispheric, perimesencephalic and intraventricular (p = 0.019), and PICA and perimesencephalic and intraventricular (p = 0.019), and PICA and perimesencephalic and intraventricular (p = 0.019), and PICA and perimesencephalic and intraventricular (p = 0.019), and PICA and perimesencephalic and intraventricular (p = 0.019), and PICA and perimesencephalic and intraventricular (p = 0.019), and PICA and perimesencephalic and intraventricular (p = 0.019), and PICA and perimesencephalic and intraventricular (p = 0.019), and PICA and perimesencephalic and intraventricular (p = 0.019), and PICA and perimesencephalic and intraventricular (p = 0.019), and PICA and perimesencephalic and intraventricular (p = 0.019).

**Conclusions:** Regional automatised volumetry seems a reliable and objective tool to quantify and describe the distribution of blood within the subarachnoid spaces. This tool accurately predicts the location of the ruptured aneurysm; its use may be prospectively considered in the emergency setting when speed and simplicity are attained.

1130-1473/© 2023 Sociedad Española de Neurocirugía. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.